

Article

## Improving Complex Treatment by Determining Sensitivity to Antibiotics During Antibacterial Therapy in Facial Surgery

Khaydarova Nargiza Mukhiddinovna

Email address: khaydarova.11011982@gmail.com  
Assistant, Bukhara State Medical University,  
Republic of Uzbekistan, city of Bukhara

**ANNOTATION.** Antibacterial therapy is the most common and effective treatment for many diseases in the field of facial surgery, including facial abscess and cellulitis. The effectiveness of the method in dental practice is discussed. The studies were conducted on 55 patients aged 18 to 45 years. of which 39 men (70.9%), 16 women (29.1%) Patients of the clinic of the Bukhara Medical Institute "Maxillofacial surgery and "reconstructive plastic surgery"" in February-May 2024 conducted a study to determine sensitivity to antibiotics .

**Key words:** antibacterial therapy, antibiotic, sensitivity, complex treatment, disc diffusion method.

**Citation:** Alkhafagi, R. K. A., Oudah, I. H., & Hillawi, R. K. A. Evaluating Health Outcomes for Iraqi Patients with Glaucoma. *Procedia of Engineering and Medical Sciences* 2024, 9(3), 110-112.

Received: 8<sup>th</sup> June 2024  
Revised: 15<sup>th</sup> June 2024  
Accepted: 22<sup>nd</sup> June 2024  
Published: 29<sup>th</sup> June 2024



**Copyright:** © 2024 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

### 1. Introduction

**Introduction:** Currently, facial surgeons are faced with the situation of "prescribing antibiotics to patients" when treating every patient. Antibacterial therapy is the most common and effective method of treating many diseases in the field of facial surgery. This is due to the high efficiency and relative harmlessness of the method. Analysis of drugs to determine sensitivity to antibiotics improves complex treatment of patients with maxillofacial surgery. Antibiotic sensitivity testing is used not only in facial surgery, but also in the treatment of many other diseases. It is better to begin treatment of any disease for which antibacterial therapy is used by determining the sensitivity of pathogenic microorganisms to antibiotics.

Modern pharmaceuticals have provided medical workers with a wide selection of antibacterial drugs. Often more than three different drugs are used to treat the same pathogen.

However, it is not always possible to immediately choose the right medicine. This is due to the presence of concomitant diseases in patients and the impossibility of taking medications individually. Like any laboratory test, an antibiotic sensitivity test is completely safe and does not cause discomfort or pain. A sample of biomaterial is taken from the patient and then tested in the laboratory.

**Purpose of the study:** analysis to determine sensitivity to antibiotics, achieving faster and safer treatment of patients by choosing the most effective drug in each individual case.

**Materials and methods:** The study was conducted on 55 patients aged 18 to 45 years, of which 39 men (70.9%), 16 women (29.1%) in the clinic of the Bukhara Medical Institute "Maxillofacial surgery and reconstructive plastic surgery" ". In the months of February-May 2024, in order to determine sensitivity to antibiotics, it was carried out using the "diffusion disk" method.

Results of the study: The study was conducted on 55 patients aged 18 to 45 years, of which 39 men (70.9%), 16 women (29.1%) in the clinic of the Bukhara Medical Institute "Maxillofacial surgery and reconstructive plastic surgery, A". The study was conducted in February-May 2024 to determine sensitivity to antibiotics and was conditionally divided into 2 groups: 1 - main group, consisting of 35 patients (25 men, 10 women), and 2 control groups - 20 people each. (12 men, 8 women).

In the main group of patients, the disk diffusion method was used to determine sensitivity to antibiotics. In this method, a bacterial suspension of a specified density (usually equal to the McFarland turbidity standard of 0.5) is applied to determine sensitivity. the surface of the agar in a Petri dish, and then add a certain number of disks containing antibiotics.

Diffusion of the antibiotic into the agar leads to the formation of a zone around the disks that inhibits the growth of microorganisms. After overnight incubation of the dishes in a thermostat at a temperature of 35 o -37 o C, the result is calculated by measuring the diameter of the zone around the disk in millimeters. The disk method was qualitative and made it possible to assess the sensitivity or resistance of the microbe to the drug.

Conclusion: antibacterial therapy in facial surgery by determining the sensitivity of bacteria to antibiotics determines the following effective results in patients; determine the most effective drug, exclude the "attachment" of pathogenic microorganisms to therapy, replace the drug due to its ineffectiveness or allergic reactions; improving the patient's social status by reducing the treatment period

#### LITERATURES:

1. Haydarova N.M. ENHANCING THE ENCOMPASSING REHABILITATION OF INDIVIDUALS INFLUENCED BY LOWER JAW DISORDERS *Journal of Healthcare and Life-Science Research* Vol. 2, No. 11, 2023 ISSN:2181-4368, 70-73
2. Haydarova N.M. Improving the Entire Rehabilitation of People Impacted with Mandible Problems *Miasto Przyszłości Kielce* 2023 Impact Factor: 9.2 ISSN-L: 2544-980X, 354-357
3. Haydarova N.M. Increasing Rehabilitation Measures for Patients Who Have Maxillofacial Thermal Wounds, *Miasto Przyszłości Kielce* 2023 Impact Factor: 9.9 ISSN-L: 2544-980X,30-32
4. Haydarova N.M. RAISING REHABILITATION METHODS FOR PATIENTS WITH MAXILLOFACIAL THERMAL INJURIES, *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE* Vol. 4 No. 2 (Feb - 2024) ISSN: 2795-921X, 422-424
5. Haydarova N.M. СОВЕРШЕНСТВОВАНИЕ РЕАБИЛИТАЦИОННЫХ МЕР У ПАЦИЕНТОВ С ТЕРМИЧЕСКИМИ ОЖОГАМИ ЧЕЛЮСТОЛИЦЕВОЙ ОБЛАСТ, *AMALIY VA TIBBIYOT FANLARI ILMIIY JURNALI*, ISSN: 2181-3469 Jild: 03 Nashr:02 2024 yil, 322-325
6. SA Kambarova EFFECT OF SURGICAL MANIPULATION TO MORPHOMETRIC DEVELOPMENT OF FACE AND JAW IN PATIENTS WITH CONGENITAL LIP AND PALATE SPLITS // *Новый день в медицине*, 2021- P. 128 - 130. 2.
7. SA Kambarova Effect of Surgical Manipulation in Morphometric Growth of Maxillofacial Area at Children with Congenital Lip and Palate Splits At I and Ii Period of Childhood// *Annals of the Romanian Society for Cell Biology*, 1853-1858. – 2021. - Vol. 25. - Issue 4. – P. 1853 – 1858.
8. KS Alixuseynovna Identification of the morphometric parameters of the cranio-fascial region of children with congenital cleft and palate reflections using a developed research map // *Central Asian Journal of Medical and Natural Science* 2 (3), 286-290 Vol. 2. - Issue 3. – P. 286 – 290. 4.
9. ША Камбарова, ШК Пулатова REVITALIZATION OF NONSPECIFIC IMMUNITY FACTORS IN PATIENTS WITH DIFFUSE PHLEGMON OF THE MAXILLOFACIAL AREA USING A BAKTERIOPHAGE // *Новый день в медицине*, 128-130 // *New day in medicine*. - 2020. - P. 128 - 130. 5.
10. KSA Xuseynovna Optimization of the Diagnosis and Treatment of Oral Epulis Based on Morphological and Cytological Analysis // *Texas Journal of Medical Science* 6, 24-26 6.
11. KS Alikhuseynovna Statistical Processing Of Morphometric Measurements Of Craniofacial Area Of Children With Congenital Cleft Labia And Palate I And II Of The Childhood Period // *Zien Journal of Social Sciences and Humanities* 5, 31-35 7.
12. SA Kambarova, GS Yadgarova CHARACTERISTIC OF MORPHOMETRIC PARAMETERS OF CRANIOFASCIAL REGION OF CHILDREN WITH CONGENITAL CLEFT LIP AND PALATE // *Academic research in educational*

---

sciences 2 (9), 295-303 8.

13. KS Alixuseynovna EFFECT OF SURGICAL MANIPULATION TO MORPHOMETRIC DEVELOPMENT OF FACE AND JAW IN PATIENTS WITH CONGENITAL LIP AND PALATE SPLITS // Web of Scientist: International Scientific Research Journal 2 (09), 29-35
14. Арипова Н., Ганиева Н. А. Clinical and prognostic characteristics of " disease-specific" autoantibodies in systemic scleroderma. – 2021.
15. Зияева Ф. К. и др. Особенности поражения нервной системы при анкилозирующем спондилоартрите : дис. – Ташкент, 2023.